

Title: Complexes of copper(II) with 3-(ortho-substituted phenylhydrazo)pentane-2,4-diones: syntheses, properties and catalytic activity for cyclohexane oxidation

Author(s): Kopylovich, Maximilian N.¹; Nunes, Andreia C. C.¹; Mahmudov, Kamran T.¹; Haukka, Matti²; Mac Leod, Tatiana C. O.¹; **Martins, Luisa M. D. R. S.**^{1,3}; Kuznetsov, Maxim L.¹; Pombeiro, Armando J. L.¹

Source: Dalton Transactions

Volume: 40 **Issue:** 12 **Pages:** 2822-2836 **DOI:** : 10.1039/c0dt01527j **Published:** 2011

Document Type: Article

Language: English

Abstract: Reactions of copper(II) with 3-phenylhydrazopentane-2,4-diones $X-2-C_6H_4-NHN = C\{C(=O)CH_3\}_2$ bearing a substituent in the ortho-position [$X = OH$ (H2L1) 1, AsO_3H_2 (H3L2) 2, Cl (HL3) 3, SO_3H (H2L4) 4, $COOCH_3$ (HL5) 5, $COOH$ (H2L6) 6, NO_2 (HL7) 7 or H (HL8) 8] lead to a variety of complexes including the monomeric $[CuL_4(H_2O)_2] \cdot H_2O$ 10, $[CuL_4(H_2O)_2]$ 11 and $[Cu(HL_4)_2(H_2O)_4]$ 12, the dimeric $[Cu_2(H_2O)_2(\mu-HL_2)_2]$ 9 and the polymeric $[Cu(\mu-L-6)](n)$ 13 ones, often bearing two fused six-membered metallacycles. Complexes 10-12 can interconvert, depending on pH and temperature, whereas the Cu(II) reactions with 4 in the presence of cyanoguanidine or imidazole (im) afford the monomeric compound $[Cu(H_2O)_4\{NCNC(NH_2)_2\}_2](HL_4)_2 \cdot 6H_2O$ 14 and the heteroligand polymer $[Cu(\mu-L-4)(im)](n)$ 15, respectively. The compounds were characterized by single crystal X-ray diffraction (complexes), electrochemical and thermogravimetric studies, as well as elemental analysis, IR, 1H - and ^{13}C -NMR spectroscopies (diones) and ESI-MS. The effects of the substituents in 1-8 on the HOMO-LUMO gap and the relative stability of the model compounds $[Cu(OH)(L-8)(H_2O)] \cdot H_2O$, $[Cu(L-1)(H_2O)_2] \cdot H_2O$ and $[Cu(L-4)(H_2O)_2] \cdot H_2O$ are discussed on the basis of DFT calculations that show the stabilization follows the order: two fused 6-membered > two fused 6-membered/5-membered > one 6-membered metallacycles. Complexes 9, 10, 12 and 13 act as catalyst precursors for the peroxidative oxidation (with H_2O_2) of cyclohexane to cyclohexanol and cyclohexanone, in MeCN/ H_2O (total yields of ca. 20% with TONs up to 566), under mild conditions.

KeyWords Plus: Mild Peroxidative oxidation; O-2- H_2O_2 -Vanadium Derivative-Pyrazine-2-Carboxylic Acid; Hydrogen-Peroxide; Crystal-Structures; Beta-Diketones; Triethanolamine Complexes; Potential Antineoplastics; Hydrocarbon Oxygenations; Azo Derivatives, Alkanes

Reprint Address: Pombeiro, AJL (reprint author), Univ Tecn Lisboa, Ctr Quim Estrutural, Inst Super Tecn, Complexo 1, Av Rovisco Pais, P-1049001 Lisbon, Portugal.

Addresses:

1. Univ Tecn Lisboa, Ctr Quim Estrutural, Inst Super Tecn, P-1049001 Lisbon, Portugal
2. Univ Joensuu, Dept Chem, FIN-80101 Joensuu, Finland
3. ISEL, Area Dept Engn Quim, P-1059007 Lisbon, Portugal

E-mail Address: pombeiro@ist.utl.pt

Funding:

Funding Agency	Grant Number
Foundation for Science and Technology (FCT), Portugal	
FEDER	

Publisher: Royal Soc Chemistry

Publisher Address: Thomas Graham House, Science Park, Milton RD, Cambridge CB4 0WF, Cambs, England

ISSN: 1477-9226

Citation: KOPYLOVICH, Maximilian N.; NUNES, Andreia C. C.; MAHMUDOV, Kamran T.; HAUKKA, Matti; MAC LEOD, Tatiana C. O.; MARTINS, Luisa M. D. R. S.; KUZNETSOV, Maxim L.; POMBEIRO, Armando J. L. - Complexes of copper(II) with 3-(ortho-substituted phenylhydrazo)pentane-2,4-diones: syntheses, properties and catalytic activity for cyclohexane oxidation. Dalton Transactions. Vol. 40, n.º 12 (2011) p. 2822-2836.